

# Journal of Applied Crystallography

Volume 41 Issue 1 Page 27-30, February 2008

**To cite this article:** Rajani K Boruah, Binoy K Saikia, Bimala P Baruah, Nibaran C Dey (2008) X-ray scattering study of the average polycyclic aromatic unit in Ledo coal  
Journal of Applied Crystallography 41 (1) , 27–30 doi:10.1107/S0021889807049655

[Prev Article](#) [Next Article](#)

**Welcome to Blackwell Synergy - the source of highly cited peer-reviewed society journals from Blackwell Publishing**

You are attempting to access the PDF of this article. To access this content you, or your library, will need to have an online subscription to the journal. Alternatively you can purchase immediate access to the article using a credit card.

- If you **already have an online subscription** please login at the top of this page.
- To **purchase an online subscription** to this journal please visit the [journal homepage](#) and click on 'Subscribe'.
- To **purchase immediate access to this article for 30 days** through our secure web site using a credit card, please click the 'Full Text Article' or 'PDF' button below, and follow the instructions.

Abstract

## X-ray scattering study of the average polycyclic aromatic unit in Ledo coal

- Rajani K.Boruah ,
- Binoy K.Saikia ,
- Bimala P.Baruah and
- Nibaran C.Dey

Rajani K. Boruah, e-mail: [rajani\\_boruah@yahoo.co.in](mailto:rajani_boruah@yahoo.co.in)

### Abstract

This report is an attempt to understand the short-range structural features, to determine the relationship(s) between the aryl/alkyl carbon ratio, and to determine the size of the average polycyclic aromatic unit in Ledo coal from Makum coalfield, Assam, India. An X-ray scattering analysis of the average polycyclic aromatic unit in the coal indicates that the aromatic fraction in this coal is 74% with the aliphatic fraction correspondingly estimated to be 26%. The average carbon atom has 2.3 nearest carbon atom neighbours at an average bond distance of 1.50 (1) Å. The average stacking height of the parallel aromatic layers and the average diameter of the aromatic layers are estimated to be 7.58 Å and 4.86 Å, respectively. For this coal, the average number of stacking layers and the average number of atoms per layer are estimated to be 2 and 8, respectively. In addition, the Gamma band is observed at a  $d$  value of 4.42 Å.