

4. Schriever, J. R. and Congalton, R. G., Evaluating seasonal variability as an aid to cover-type mapping from Landsat Thematic Mapper data in the northeast. *Photogramm. Eng. Remote Sensing*, 1995, **61**, 321–327.
5. Watson, N. and Wilcock, D., Preclassification as an aid to the improvement of thematic and spatial accuracy in land cover maps derived from satellite imagery. *Remote Sensing Environ.*, 2001, **75**, 267–278.
6. Asner, P. G., Biophysical and biochemical sources of variability in canopy reflectance. *Remote Sensing Environ.*, 1998, **64**, 234–253.
7. Kumar, L., Schmidt, K., Dury, S. and Skidmore, A., Imaging spectrometry and vegetation science. In *Imaging Spectrometry, Basic Principles and Prospective Applications* (ed. Van Der Meer, F. D.), Kluwer, 2001, p. 111.
8. Lewis, M. M. and Dunk, I., Seagrass and shallow water feature discrimination using HyMap. In Proceedings of the Tenth Australasian Remote Sensing and Photogrammetry Conference, Adelaide, Australia, 21–25 August 2000, vol. 1, pp. 1092–1108 (CD-ROM).
9. Shippert, P., Introduction to hyperspectral image analysis. Ph D thesis, Geography Department, University of Auckland, New Zealand, 1992, pp. 492–504.
10. Clark, R. N. and Swayze, G. A., Mapping minerals, amorphous materials, environmental materials, vegetation, water, ice and snow, and other materials: The USGS tricorder algorithm. In *Summaries of the Fifth Annual JPL Airborne Earth Science Workshop*, JPL Publication, 1995, 95-1, vol. 1, pp. 39–40.
11. Aber, J. D. and Martin, M. E., High spectral resolution remote sensing of canopy chemistry. In *Summaries of the Fifth JPL Airborne Earth Science Workshop*, JPL Publication, 1995, 95–1, vol. 1, pp. 1–4.
12. Merton, R. N., Multi-temporal analysis of community scale vegetation stress with imaging spectroscopy. Ph D thesis, Geography Department, University of Auckland, New Zealand, 1992, p. 492.
13. Pearlman, J. S., Barry, P. S., Segal, C. C., Shepanski, J., Beiso, D. and Carman, S. L., Hyperion, a space-based imaging spectrometer. *IEEE Trans. Geosci. Remote Sensing*, 2003, **41**, 1150–1173.
14. Landgrebe, D., Some fundamentals and methods for hyperspectral image data analysis. Presented at workshop SPIE Photonics West, San Jose, CA, 1999, pp. 1–10.
15. Woodcock, C. E., Collins, J. B., Gopal, S., Jakabhazy, V. D., Harward, V. J., Levitan, J. and Wu, Y., Mapping forest vegetation using Landsat TM imagery and a canopy reflectance model. *Remote Sensing Environ.*, 1994, **50**, 240–254.
16. Gilbert, M. A., Garcia-Haro, F. J. and Melia, J., A mixture modeling approach to estimate vegetation parameters for heterogeneous canopies in remote sensing. *Remote Sensing Environ.*, 2000, **72**, 328–345.
17. Leckie, D. G., Gougeon, F. A., Walsworth, N. and Paradine, D., Stand delineation and composition estimation using semi-automated individual tree crown analysis. *Remote Sensing Environ.*, 2003, **85**, 355–369.
18. Sabnis, S. D. and Amin, J. C., Eco-environmental studies of Sardar Sarovar environs. The M.S. University of Baroda, 1992.
19. ImSpec, The ACORN home page, 2004; <http://www.imspec.com/index.html>
20. Leica, ATCOR for Imagine, Data Sheet, 2004; <http://www.geosystems.de/atcor/index.html>
21. Shih, T. Y., On the atmospheric correction for a Hyperion scene. Presented in 25th Asian Conference and 1st Asian Space Conference on Remote Sensing, Thailand, 22–24 November 2004.
22. Research Systems, Inc, ENVI Version 3.6, 2002; <http://www.RSInc.com/envi>
23. Green, A. A., Berman, M., Switzer, P. and Craig, M. D., A transformation for ordering multispectral data in terms of image quality with implications for noise removal. *IEEE Trans. Geosci. Remote Sensing*, 1988, **26**, 65–74.
24. Datt, B., McVicar, T. R., Jupp, D. L. B. and Pearlman, J. S., Pre-processing EO-1 hyperion hyperspectral data to support the application of agricultural indexes. *IEEE Trans. Geosci. Remote Sensing*, 2003, **41**, 1246–1259.
25. Jupp, D. P. L., Datt, B., Lovell, J. and King, E., EO-1/hyperion data workshop notes. CSIRO Office of Space Science and Applications, Earth Observation Centre, Canberra, Australia, April 2002.
26. Clark, R. N., Spectroscopy of rocks and minerals, and principles of spectroscopy. In *Manual of Remote Sensing, Vol. 3, Remote Sensing for the Earth Sciences* (ed. Rencz, A. N.), John Wiley, New York, 1999, p. 137.
27. Robilla, S. A., Using spectral distances for speedup in hyperspectral image processing. *Int. J. Remote Sensing*, 2005, **26**, 5629–5650.
28. Price, J. C., How unique are spectral signatures? *Remote Sensing Environ.*, 1994, **49**, 181–186.

ACKNOWLEDGEMENTS. We are grateful for the funding support by Indian Space Research Organization, Bangalore. We are thankful to the Forest Department, Gujarat for logistic support.

Received 9 January 2007; revised accepted 17 September 2007

Dinitrogen fixation activity of *Azospirillum brasilense* in maize (*Zea mays*)

S. P. Saikia^{1,2,*}, Vanita Jain¹, Sangeeta Khetarpal¹ and Samitha Aravind¹

¹Division of Plant Physiology, Indian Agricultural Research Institute, New Delhi 110 012, India

²Division of Medicinal, Aromatic and Economic Plants, North East Institute of Science and Technology (CSIR), Formerly Regional Research Laboratory, Jorhat 785 006, India

The association of maize with the N₂-fixing bacteria, *Azospirillum* was examined to evaluate possible benefits for agriculture. Maize seedlings develop nodule-like tumour knots (*para*-nodules) along primary roots when treated with the auxin 2,4-dichlorophenoxyacetic acid (2,4-D). The presence of nitrogenase activity (acetylene reduction assay) was noticed in plants treated with *Azospirillum* along with 2,4-D (nodulated plants) at various stages of growth. The 2,4-D induced nodule-like structures provided niche for the establishment of the bacteria, thereby enhancing nitrogen fixation in the nodulated roots. *Azospirillum* inoculation had a significantly positive effect on the activity of the enzymes glutamine synthetase and glutamate synthase in leaves of *para*-nodulated maize plants. The *para*-nodulated plants showed higher rates of net photosynthesis, stomatal conductance leading to higher yield compared to non-nodulated (control) plants. The

*For correspondence. (e-mail: sp_sidhu@yahoo.com)

nutrient content in grains and stover was higher in the case of *para*-nodulated plants. The results provide evidence that maize plants are potentially able to create a symbiosis with diazotrophic bacteria, which colonize *para*-nodule tissue intracellularly, promoting a higher level of N₂ fixation for better growth and development of plants. Further, our data suggest that an increased understanding of plant root–rhizosphere N₂-fixing bacterial interactions is required before any consistent, significant and beneficial N₂-fixing association can be developed.

Keywords: Acetylene reduction, *Azospirillum brasilense*, maize, physiological activities, seedling inoculation.

The times have changed; the face of the country is changed; the quality of the soil has changed; and if we will live as well, and become as rich and respectable as our fathers, we must cultivate their virtues; but abandon their farming system.

The Farmers Manual (1819)

BIOLOGICAL nitrogen fixation is the key to sustain agricultural productivity through the application of biofertilizers in the field. Increased foodgrain cannot be produced unless we carefully make use of biological nitrogen fixation. Since the nodulation per se is a physiological process, it is possible to induce nodulation using different plant growth regulators and it is obvious that the nitrogen-fixing bacteria can enter through cracked epidermal cells of the nodules and such nodules can provide a niche for the bacteria that fixes nitrogen inside the nodules¹. Among different plant hormones, 2,4-dichlorophenoxyacetic acid (2,4-D), a synthetic auxin, is the best in inducing nodular outgrowth in cereals². The objectives of this study were to determine the amount of N₂-fixation by *Azospirillum* in association with *Zea mays*, and to measure whether inoculation with this organism established an association that could contribute significant amounts of fixed N to plants.

Seeds of *Z. mays* L., cultivar Kiran were surface-sterilized and germinated on sterile, acid-washed gravel culture at 25°C for 5–6 days. Uncontaminated seedlings from the acid-washed gravel culture were grown in glass tubes (20 cm length, 3 cm diameter, one seedling per tube) containing sterile N-free Hoagland solution at 25°C in growth room. Exposure of the root zone to light was prevented. Bacterial inoculations were performed two days after transferring the seedlings to Hoagland solution with 0.1 ml of *Azospirillum brasilense* (Sp-29, culture collection, Division of Microbiology, IARI, New Delhi) culture grown at room temperature for 24 h, containing 10⁷–10⁸ cells/ml along with 1.0 ppm 2,4-D. Nodule-like structure were formed on the roots after a period of 7–10 days after treatment with 2,4-D. The two-week-old seedlings after nodulation under laboratory conditions were transferred to pots (50 cm × 50 cm × 50 cm) in the net house under

natural conditions. Each pot was filled with 45 kg of soil. After transplanting in pots, biofertilizer *Azospirillum* was applied in the form of solution 20 ml/pot @ 10⁷–10⁸ cells/ml. No chemical fertilizers were applied. Nitrogenase activity was assayed in actively growing intact nodulated roots at 30, 60 and 90 days after transplanting (DAT) in pots following the acetylene reduction assay (ARA) method³, with slight modifications: 5 cc of gas or air was removed and 5 cc of acetylene was injected into the vials containing 1 g of nodulated roots and incubated for 12 h at 30°C. Gas mixture (2 cc) was fed to the gas chromatograph and the acetylene reduced was measured quantitatively. Glutamine synthetase (GS) and glutamate synthase (GOGAT) activities were assayed in actively growing leaves of *para*-nodulated plants at 30, 60 and 90 DAT. The activities of GS and GOGAT were estimated using the spectrophotometric method⁴. Two grams of leaves was ground in extraction buffer, 100 mM Tris buffer (pH 7.5). The homogenate was filtered through a cheese cloth and centrifuged at 10,000 rpm at 4°C for 15 min. The assay for GS was carried out at 37°C (Reaction mixture consisted of Tris, 100 μmol; MgSO₄, 100 μmol; cysteine, 10 μmol; L-glutamate, 250 μmol; ATP, 10 μmol; enzyme extract, 0.1–0.5 ml, and hydroxylamine, 20 μmol in the reaction volume of 3 ml). The reaction was terminated by adding ferric chloride reagent

Table 1. Nitrogenase activity (nmol of ethylene/g fr. wt/h) in roots of control and *para*-nodulated maize plants

Treatment	Days after transplanting		
	30	60	90
Control	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00
Nodulated	1.13* ± 0.34	2.68* ± 0.36	2.86* ± 0.42

*Values were significant at 5% level of probability.

Table 2. Glutamine synthetase activity (μmol/g fr. wt/h) in leaves of control and *para*-nodulated maize plants

Treatment		Days after transplanting		
		30	60	90
Control	Leaf	89.57 ± 0.42	148.80 ± 1.40	139.33 ± 3.64
Nodulated	Leaf	102.64* ± 1.73	188.27* ± 1.43	167.59* ± 2.48

*Values were significant at 5% level of probability.

Table 3. Glutamate synthase activity (μmol NADH oxi/min/g fr. wt) in leaves of control and *para*-nodulated maize plants

Treatment		Days after transplanting		
		30	60	90
Control	Leaf	37.87 ± 1.09	100.80 ± 0.00	55.95 ± 0.00
Nodulated	Leaf	25.11 ± 1.03	132.59* ± 3.12	84.78* ± 1.10

*Values were significant at 5% level of probability.

RESEARCH COMMUNICATIONS

Table 4. Photosynthetic rate ($\mu\text{mole CO}_2/\text{sq. m/S}$) and stomatal conductance (m/S) in control and *para*-nodulated maize plants

Days after transplanting	Treatment			
	Photosynthetic rate		Stomatal conductance	
	Control	Nodulated	Control	Nodulated
30	3.70 \pm 0.38	11.17* \pm 0.06	0.06 \pm 0.01	0.12* \pm 0.06
60	5.40 \pm 0.77	12.98* \pm 0.18	0.16 \pm 0.05	0.32* \pm 0.08
90	6.90 \pm 0.20	14.28* \pm 0.13	0.22 \pm 0.03	0.36* \pm 0.07

*Values were significant at 5% level of probability.

Table 5. Fresh and dry weight of leaf (g/plant) and leaf area (sq. cm) in control and *para*-nodulated maize plants

Days after transplanting	Treatment	Leaf		
		Fresh weight	Dry weight	Leaf area
30	Control	23.99 \pm 1.94	5.90 \pm 0.32	1527.84 \pm 41.91
	Nodulated	42.63* \pm 2.34	7.86* \pm 0.51	1647.89* \pm 89.45
60	Control	35.96 \pm 0.41	7.01 \pm 0.17	1838.24 \pm 31.50
	Nodulated	46.43* \pm 0.21	9.49* \pm 0.13	2187.21* \pm 64.04
90	Control	20.65 \pm 1.54	6.85 \pm 0.20	1286.83 \pm 47.97
	Nodulated	32.73* \pm 0.65	11.54* \pm 0.87	1562.29* \pm 40.34

*Values were significant at 5% level of probability.

Table 6. Fresh and dry weight of stem (g/plant) in control and *para*-nodulated maize plants

Days after transplanting	Treatment	Stem	
		Fresh weight	Dry weight
30	Control	34.88 \pm 0.39	3.14 \pm 0.04
	Nodulated	44.53* \pm 2.57	7.40* \pm 0.55
60	Control	75.99 \pm 1.72	15.33 \pm 0.58
	Nodulated	101.22* \pm 5.71	22.80* \pm 1.83
90	Control	49.94 \pm 1.68	17.42 \pm 0.93
	Nodulated	112.57* \pm 2.57	39.70* \pm 1.92

*Values were significant at 5% level of probability.

Table 7. Yield and yield components in *para*-nodulated maize at final harvest

Treatment	Ear length (cm)	No. of grains per ear	Hundred grain weight (g)
Control	13.33 \pm 0.08	248.67 \pm 1.02	22.07 \pm 0.02
Nodulated	16.50* \pm 0.09	329.67* \pm 2.03	32.39* \pm 0.02

*Values were significant at 5% level of probability.

and absorbance taken at 540 nm. The reaction mixture for GOGAT contained Tris, 75 μmol , α -ketoglutaric acid, 10 μmol ; L-glutamine, 15 μmol , NADH, 0.3 μmol , and enzyme aliquot, 100 μL in the total reaction volume of 3 ml. Readings were taken at 340 nm (UV-Vis spectrophotometer, EC).

The fresh and dry weight of the stem and leaf and leaf area was measured at 30, 60 and 90 DAT. The stem dry

weight was measured after drying the material for 72 h at 70°C. The leaf dry weight was determined after drying the leaves to a constant weight and leaf area was measured using a leaf area meter (LI-3100, LI-COR, Inc., Lincoln, NE, USA). Photosynthetic rate and stomatal conductance were measured with the help of an Infra Red Gas Analyser (LI-COR 6200). The yield components as well as the nitrogen, phosphorus and potassium content in stover and grain were estimated at final harvest. Dried plant samples were ground to a fine powder and used for the estimation of nutrients. Nitrogen was estimated using an auto analyser (Technicon, USA). Phosphorus was estimated using standard procedure⁵ and potassium in a flame photometer (model CL 22D). For estimation of phosphorus and potassium, samples were digested with the diacid mixture. The experiment was laid out in completely randomized design. All the observations are the means of three replicates and comparisons of treatment means were made at the 5% confidence level⁶.

The 2,4-D treatment (1.0 ppm) induced the nodule-like tumour knots (*para*-nodules) in maize roots. It started from the steler portion, and cell division started at the inner cortex level and pushed the epidermal cells. These cells bulge out in the shape of a nodule. The inoculated bacteria *Azospirillum* entered through loosening of the epidermis without causing much damage to infected host epidermal cells^{7,8}. Acetylene reduction rate was observed in nodulated plants inoculated with *Azospirillum brasilense* at different stages of growth. Control seedlings showed no acetylene reduction (Table 1). Nitrogenase ac-

Table 8. NPK content (%) in grain and stover in *para*-nodulated maize at final harvest

Treatment	N content		P content		K content	
	Grain	Stover	Grain	Stover	Grain	Stover
Control	1.39 ± 0.08	0.94 ± 0.04	0.11 ± 0.02	0.09 ± 0.04	0.10 ± 0.02	0.28 ± 0.06
Nodulated	1.52* ± 0.09	1.24* ± 0.06	0.13* ± 0.03	0.11* ± 0.02	0.11* ± 0.02	0.30* ± 0.07

*Values were significant at 5% level of probability.

tivity of *A. brasilense*, in terms of associative symbiotic nitrogen fixation, has been reported by many workers^{9,10}. The 2,4-D induced nodule-like structure serve as a niche for bacterial colonization in maize roots, thereby enhancing the nitrogen-fixing capacity in the roots. Enhanced nitrogenase activity observed in maize *para*-nodules might be due to the efficiency of the host to encapsulate the intracellular colonizing bacterial cells with a membrane layer resembling the peri-bacteroid membrane of the legume nodule². This also indicates that the host plant was capable of supplying the necessary substrate for N₂-fixation (C₂H₂ reduction) of the bacteria residing mainly in the *para*-nodule. Biological nitrogen fixation during nodular association or *nif* gene transfer to the plants is high¹¹. The bacteria better express their N₂-fixing potential inside nodular tissues due to lower competition for nutrients and protection against high level of O₂ pressure on the root surface¹². Demonstration of significant N₂-fixation in maize seedling roots using 2,4-D and *Azospirillum* is an important development, which may open new fields for both basic and applied aspects of N₂-fixation.

Plants treated with *Azospirillum* along with 2,4-D (nodulated plants) showed increased activity of GS and GOGAT in leaves (Tables 2 and 3), and increased photosynthetic rate and stomatal conductance when compared with uninoculated controls at different stages of growth (Table 4). These results are in accordance with the earlier findings reported in wheat plants¹³. The fresh and dry weight of leaf and stem (Tables 5 and 6) were also found to be higher in the nodulated plants at different stages of growth. The higher biomass production in nodulated plants may be due to higher leaf area (Table 5) and better root and shoot growth, which tends to absorb more water and nutrients from the deeper zone of the soil combined with the better nitrogen fixation capacity of the *Azospirillum*-treated nodulated plants¹⁴.

Data pertaining to the yield parameters (Table 7) showed that inoculation with *A. brasilense* along with 2,4-D produced higher number of ears per plant with increased ear length, ear weight and higher number of grains per ear. Yield increases obtained in inoculated plants, however, have been attributed to biological nitrogen fixation and may also be due to the production of growth substance by the colonizing bacteria. Increased N supply through N₂ fixation, higher nutrient uptake¹⁵ and higher

photosynthetic rate and stomatal conductance in inoculated plants might have contributed to the increase in grain yield.

In the present experiment no chemical source of N was added and the significant increase in the level of nutrients was obtained. The major nutrients, viz. nitrogen, phosphorus and potassium in grains and stover increased significantly in *para*-nodulated maize plants compared to control (Table 8). Data on NPK content in grains and stover at final harvest in plants revealed that while nitrogen and phosphorus content were higher in grains, potassium was high in stover. Increase in nitrogen content in shoot and grains has been reported in cereal¹⁶. Increased uptake of mineral ion by maize and sorghum has been reported under laboratory conditions¹⁷.

Proper understanding of plant bacterial associations is necessary if we are to increase the N nutrient input by rhizosphere bacteria and improve the N nutrition of cereal crops by this method. The population dynamics of introduced *A. brasilense* inside *para*-nodules must be studied during the entire period of plant growth and the overall N gain of the symbiotic system must be investigated by N-balance calculations or with ¹⁵N-isotope techniques. Studies of associative symbiosis should be pursued vigorously because of their potential importance, and because the plant and bacterial association examined to date has been limited.

1. Ridge, R. W., Bander, G. B. and Rolfe, B. G., Nodule-like structures induced on the roots of wheat seedlings by the addition of synthetic auxin 2,4-D and the effects of microorganisms. *Aust. J. Plant Physiol.*, 1992, **19**, 481–492.
2. Christiansen Weniger, C., Ammonium excreting *Azospirillum brasilense* C3: GUSA inhabiting induced tumors along stem and roots of rice. *Soil Biol. Biochem.*, 1997, **29**, 943–950.
3. Hardy, R. W. F., Hilsten, R. D., Jackson, E. K. and Burns, R. C., The acetylene assay for nitrogen fixation: Laboratory and field evaluation. *Plant Physiol.*, 1968, **43**, 1195–1207.
4. Mohanty, B. and Fletcher, J. S., Ammonium influence on nitrogen assimilatory enzymes and protein accumulation in suspension cultures of pearl scarlet rose. *Physiol. Plant.*, 1980, **48**, 453–459.
5. Jackson, M. L., *Soil Chemical Analysis*, Prentice-Hall, New Jersey, 1973.
6. Panse, V. G. and Shukhatme, P. V., *Statistical Procedures for Agricultural Workers*, ICAR Publication, New Delhi, 1967.
7. Christiansen Weniger, C. and Vanderleyden, J., Ammonium-excreting *Azospirillum* sp. become intracellularly established in maize (*Zea mays*) *para*-nodules. *Biol. Fertil. Soils*, 1994, **17**, 1–8.

8. Elanchezian, R. and Panwar, J. D. S., Effect of 2,4-D and *Azospirillum brasilense* on nitrogen fixation, photosynthesis and grain yield in wheat. *J. Agron. Crop Sci.*, 1997, **178**, 129–133.
9. Okon, Y., *Azospirillum* as a potential inoculant for agriculture. *Trends Biotechnol.*, 1985, **3**, 323–328.
10. Tilak, K. V. B. R. and Rao Subba, N. S., Association of *Azospirillum brasilense* with pearl millet (*Pennisetum americanum* (L) Leeke). *Biol. Fertil. Soils*, 1987, **4**, 97–102.
11. Dawe, D., The potential role of biological nitrogen fixation in meeting future demand for rice and fertilizer. In *The Quest for Nitrogen Fixation in Rice* (eds Ladha, J. K. and Reddy, P. M.), International Rice Research Institute, Manila, Philippines, 2000, pp. 1–9.
12. Boddey, R. M. and Dobereiner, J., Nitrogen fixation associated with grasses and cereals: Recent progress and perspectives for the future. *Fertil. Res.*, 1995, **42**, 241–250.
13. Panwar, J. D. S. and Elanchezian, R., Effect of 2,4-D and *Azospirillum brasilense* on growth and yield in the nodule-induced transplanted wheat. *Indian J. Plant Physiol.*, 1998, **3**, 143–146.
14. Kennedy, I. R. and Tchan, Y. T., Biological nitrogen fixation in non-leguminous field crops: Recent advances. *Plant Soil*, 1992, **141**, 93–118.
15. Subba Rao, N. S., Tilak, K. V. B. R. and Singh, C. S., Yield responses of crop to inoculation with *Azospirillum brasilense* in India. *Zbl. Mikrobiol.*, 1985, **140**, 97–102.
16. Baldani, V. L. D. and Dobereiner, J., Host-plant specificity in the infection of cereals with *Azospirillum* spp. *Soil Biol. Biochem.*, 1980, **12**, 433–439.
17. Tein, T. M., Tien, H. G., Gaskin, M. H. and Hubbell, D. H., Poly-galacturonic acid transeliminase production by *Azospirillum* species. *Can. J. Microbiol.*, 1981, **27**, 426–431.

ACKNOWLEDGEMENTS. Financial support from Council of Scientific and Industrial Research, New Delhi in the form of Junior Research Fellowship to the senior author is acknowledged. S.P.S. also thanks to the anonymous reviewers for their insightful suggestions and critical reviews of the manuscript.

Received 16 January 2006; revised accepted 17 August 2007

Enhancement of cytotoxicity and DNA binding of cisplatin in Dalton's lymphoma cells by α -tocopherol

Suniti Sarna* and Rakesh K. Bhola

Department of Zoology, Gauhati University, Guwahati 781 014, India

A combination treatment of cisplatin and α -tocopherol on Dalton's lymphoma cells *in vitro* was studied in order to examine the effect of α -tocopherol on cisplatin cytotoxicity and binding of platinum to DNA (DNA platination) or its removal from DNA. Cisplatin cytotoxicity which is measured as cell survival of Dalton's lymphoma cells was found to be enhanced by various factors, including concentration of drug,

treatment duration and presence of α -tocopherol in the medium. Tumour cells treated with increasing concentrations of cisplatin or cisplatin with α -tocopherol lower cell survival and increase DNA platination. There was a significant correlation between percentage survival of Dalton's lymphoma cells and DNA platination. The study of percentage removal of platinum has shown an inverse linear correlation between DNA damage and DNA repair. The enhanced cytotoxicity of cisplatin by α -tocopherol could be attributed to increased platinum binding to DNA, which might decrease DNA repair.

Keywords: Cisplatin, cytotoxicity, Dalton's lymphoma, DNA platination, α -tocopherol.

CISPLATIN is successfully used in the chemotherapy of a wide variety of experimental malignant tumours^{1–3} and also in the treatment of many human cancers^{4,5}. Many reports have been published describing the intercalation of cisplatin with DNA^{6–8}, which suggest that cellular DNA could be the primary target for cisplatin in its cytotoxicity. The antitumour activity of cisplatin is believed to result from the interaction of drug with DNA, which leads to the formation of different types of adducts through reaction of the bifunctional platinum compound with N⁷ atoms of the nucleobases guanine and adenine. The major adducts are intrastrand crosslinks formed by the binding of cisplatin on two neighbouring guanines, on adenine and guanine and on two guanines separated by one or more nucleobases. Other types of adducts formed are the inter-strand crosslinks on two guanines and monofunctionally bound cisplatin on guanines^{8–10}. Hyperthermia enhances the cytotoxic effect of cisplatin in both cisplatin-sensitive and cisplatin-resistant mammalian cell lines^{11–13}. The enhanced cytotoxicity may be due to increased DNA crosslinking by cisplatin, increased cellular accumulation of platinum or decreased repair of drug-induced DNA damage^{12–14}. The amount of platinum bound to DNA is also likely to be related to the sensitivity of cells to cisplatin¹⁵. Prasad and Rama¹⁶ showed that vitamin E induced both morphological differentiation and growth inhibition of murine and human neuroblastoma cells in culture at least in part by an antioxidant mechanism. Vitamin E probably inhibits DNA synthesis in tumour cells because it inhibits cell division. It may act similarly at the site on RNA or DNA where cisplatin binds¹⁷. Prasad *et al.*¹⁷ demonstrated the presence of vitamin E-binding proteins in tumour cells in culture, but the relationship of vitamin E-binding proteins to the mechanism of action of vitamin E is unknown. The association of a significant amount of radioactive vitamin E with the purified chromatin suggests that vitamin E may modulate genetic expression in mammalian cells¹⁷. The present study was undertaken to investigate the effects of vitamin E on cisplatin cytotoxicity and on binding of platinum to DNA (DNA platination) or its removal from DNA.

*For correspondence. (e-mail: sunitisarna@yahoo.com)